

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 770 615 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
18.06.2003 Bulletin 2003/25

(51) Int Cl.7: **C07D 487/04**
// (C07D487/04, 249:00,
239:00)

(21) Application number: **96307528.8**

(22) Date of filing: **16.10.1996**

(54) Process for the preparation of dihaloazolopyrimidines

Verfahren zur Herstellung von Dihalogenazolopyrimidinen

Procédé de production de dihaloazolopyrimidines

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL
PT SE**
Designated Extension States:
AL LT LV RO SI

(30) Priority: **27.10.1995 US 8022**

(43) Date of publication of application:
02.05.1997 Bulletin 1997/18

(73) Proprietors:
• **BASF AKTIENGESELLSCHAFT**
67056 Ludwigshafen (DE)
Designated Contracting States:
AT BE CH LI DE DK ES FI FR GB IE IT NL
• **American Cyanamid Company**
Madison, New Jersey 07940-0874 (US)
Designated Contracting States:
GR LU PT SE

(72) Inventors:
• **Krummel, Günther**
55578 Vendersheim (DE)
• **Stumm, Karl-Otto**
55459 Aspisheim (DE)
• **Pees, Klaus-Jürgen**
55129 Mainz (DE)
• **Liers, Peter Heinz Rudi**
55424 Münster-Sarmsheim (DE)

(74) Representative: **Langfinger, Klaus Dieter, Dr. et al**
Director Patents & Licensing,
BASF Aktiengesellschaft ZDX/P-C6
67056 Ludwigshafen (DE)

(56) References cited:
EP-A- 0 322 359 EP-A- 0 444 747
EP-A- 0 550 113 WO-A-95/11246
DE-A- 3 522 463 US-A- 3 907 799
US-A- 5 006 656

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

BACKGROUND OF THE INVENTION


[0001] Dihaloazolopyrimidines are useful as intermediates in the preparation of a variety of agrochemical and pharmaceutical compounds. In particular, 5,7-dihalo-6-aryl-1,2,4-triazolo[1,5-a]pyrimidines are key intermediates in the preparation of fungicidal triazolopyrimidine derivatives which are described in EP-A2-550113.

[0002] EP-A2-550113 describes a method for the preparation of 5,7-dihalo-6-aryl-1,2,4-triazolo[1,5-a]pyrimidines from malonic acid esters and 3-amino-1,2,4-triazole. However, that method is not entirely satisfactory because those pyrimidine compounds are obtained in low yield.

[0003] G. Fischer (Advances in Heterocyclic Chemistry, 1993, 57, 81-138) describes the formation of triazolopyrimidines from 1,3-dicarbonyl compounds and 3-amino-1,2,4-triazole, and states that refluxing in glacial acetic acid is "standard conditions". Y. Makisumi (Chem. Pharm. Bull., 1961, 9, 801-808) reports that under those conditions the condensation of diethyl malonate with 3-amino-1,2,4-triazole does not proceed. Makisumi discloses that the reaction could be carried out in the presence of sodium ethoxide in ethanol, and that the product dihydroxytriazolopyrimidine could be converted to the corresponding dichlorotriazolopyrimidine using a large excess of phosphorus oxychloride. However, Makisumi's method is not entirely satisfactory for the preparation of dihaloazolopyrimidines because a large excess of phosphorus oxychloride is required and the overall yield of the reactions starting from diethyl malonate is often low.

SUMMARY OF THE INVENTION

[0004] The present invention provides an effective and efficient process for the preparation of a dihaloazolopyrimidine having the structural formula I




(I)

wherein

- X_1 is chlorine or bromine;
 R is phenyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_6 alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, C_1 - C_4 alkoxycarbonyl, phenyl, phenoxy or benzyloxy groups, naphthyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, C_1 - C_4 alkoxycarbonyl, phenyl, phenoxy or benzyloxy groups, C_1 - C_6 alkyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy or C_1 - C_4 haloalkoxy groups, C_3 - C_8 cycloalkyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 -haloalkoxy groups, or C_2 - C_6 alkenyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy or C_1 - C_4 haloalkoxy groups;
 X is CR_1 or N;
 Y is CR_2 or N;
 Z is CR_3 or N;
 R_1 , R_2 and R_3 are each independently hydrogen or C_1 - C_6 alkyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, amino, C_1 - C_4 alkylamino or di(C_1 - C_4 alkyl)amino groups;
 which process comprises

(a) reacting (1) a malonic acid ester having the structural formula II




wherein R_8 and R_9 are each independently C_1 - C_6 alkyl, and R is as described above with (2) a heterocyclamine having the structural formula III



wherein X, Y and Z are as described above at a temperature of at least 100°C in the presence of at least one molar equivalent, relative to the malonic acid ester, of a base to form an intermediate salt,


(b) optionally acidifying said intermediate salt with aqueous acid to form a dihydroxyazolopyrimidine having the structural formula IV



wherein R, X, Y and Z are as described above, and

(c) halogenating the intermediate salt or dihydroxyazolo-pyrimidine with at least about two molar equivalents of a halogenating agent, e.g., phosphorus oxychloride, phosphorus oxybromide, phosphorus pentachloride or phosphorus pentabromide or a suitable mixture thereof at a temperature of at least about 100°C .

[0005] The present invention also provides an effective and efficient process for the preparation of a dihydroxyazolopyrimidine having the structural formula IV



wherein R, X, Y and Z are as described above. This product (IV) is produced by the above-described procedure wherein the intermediate salt is acidified; the product (IV) then may be isolated, if desired.

[0006] It is, therefore, an object of the present invention to provide an efficient new process for the preparation of dihaloazolopyrimidines.

[0007] It is another object of the present invention to provide a novel process for preparing dihydroazolopyrimidines.

[0008] Other objects and advantages of the present invention will be apparent to those skilled in the art from the following description and the appended claims.

DETAILED DESCRIPTION OF THE INVENTION



[0009] In one embodiment of the present invention, a malonic acid ester 5 represented by formula II is reacted with at least about one molar equivalent of a heterocyclamine represented by formula III, preferably in a temperature range of about 120°C to 200°C, more preferably about 150°C to 180°C, and in the presence of a base and/or solvent to form an intermediate salt. The intermediate salt is halogenated with at least about two molar equivalents of phosphorus oxychloride, phosphorus oxybromide, phosphorus pentachloride or phosphorus pentabromide, or a suitable mixture thereof, preferably in a temperature range of about 120°C to 150°C.

[0010] Advantageously, it has now been found that dihaloazolopyrimidines may be obtained in high yield and good purity by the effective and efficient process of the present invention. In contrast, dihaloazolopyrimidines are obtained in comparatively low yield when prepared according to art methods.

[0011] A further advantage of the present invention is that the inventive process may be conducted in one pot when the intermediate salt is not acidified. A one pot reaction sequence is highly desirable because it avoids the isolation of intermediate compounds and significantly reduces the amount of chemical waste produced.

[0012] The intermediate salt is prepared in the presence of added base. The base is present in an amount of at least about one molar equivalent relative to the malonic acid ester. Bases suitable for use in the process of the present invention include tertiary amines such as tri(C₂-C₆alkyl)amines, pyridine, substituted pyridines, quinoline, substituted quinolines, and ureas; alkali metal hydroxides such as sodium hydroxide and potassium hydroxide; alkaline earth metal hydroxides such as calcium hydroxide and magnesium hydroxide; alkali metal C₁-C₆alkoxides such as sodium ethoxide and potassium *tert*-butoxide; alkaline earth metal C₁-C₆alkoxides such as magnesium ethoxide; alkali metal carbonates such as sodium carbonate and potassium carbonate; and alkaline earth metal carbonates such as calcium carbonate. Preferred bases include tri(C₂-C₆alkyl)amines such as triethylamine and tributylamine, pyridine, 4-(N,N-dimethylamino)pyridine, quinoline, and N,N,N',N'-tetramethylurea with triethylamine and tributylamine being more preferred.

[0013] The intermediate salt of this invention is represented by structural formula V when prepared in the absence of added base, and structural formula VI when prepared in the presence of added base:



wherein R, X, Y and Z are as described above and "Base" represents the added base.

[0014] In a further preferred embodiment of the present invention, a solvent is present. Solvents suitable for use in the process of the present invention have a boiling point of at least about 80°C and include aromatic hydrocarbons such as mesitylene, toluene, xylenes and mixtures thereof; chlorinated aromatic hydrocarbons such as mono- and

dihalobenzenes and mixtures thereof; polynuclear aromatic hydrocarbons such as naphthalene, alkylnaphthalenes and mixtures thereof; alcohols such as butanol; and mixtures thereof. The solvent of the present invention preferably has a boiling point range of about 80°C to 220°C, more preferably about 120°C to 180°C. Mesitylene is one of the preferred solvents of the present invention.

[0015] The reaction between the malonic acid ester and the heterocyclamine is preferably performed at a pressure of about one atmosphere or higher. If the reaction includes a solvent having a boiling point (defined at normal atmospheric pressure) lower than the reaction temperature, the reaction pressure must be elevated so that the solvent boiling point is elevated to at least the reaction temperature.

[0016] In some embodiments of the inventive process, an aqueous acid is used to acidify the intermediate salt. Aqueous acids suitable for use include aqueous mineral acids such as hydro-chloric acid, hydrobromic acid and sulfuric acid, and aqueous organic acids such as trifluoroacetic acid with hydrochloric acid, hydrobromic acid, and sulfuric acid being preferred.

[0017] The halogenation reaction may comprise reacting the intermediate salt or the dihydroxyazolopyrimidine with a suitable halogenating agent under conditions that produce the desired dihaloazolopyrimidine. Any halogenating agent and conditions known in the art may be used.

[0018] Preferably, the halogenating agent and conditions are those described herein for the preferred embodiments of the present invention. Advantageously, the halogenation reaction may be conducted at atmospheric pressure or at a pressure greater than atmospheric pressure. The term "a suitable mixture thereof", as used in the specification and claims with regard to the halogenating agents described herein, is defined as a phosphorus oxychloride and phosphorus pentachloride mixture or a phosphorus oxybromide and phosphorus pentabromide mixture.

[0019] The process of the present invention is especially useful for the preparation of dihaloazolopyrimidines wherein

X₁ is chlorine;
 R is phenyl optionally substituted with one or more halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄-alkoxy, C₁-C₄haloalkoxy, phenyl, phenoxy or benzyloxy groups, or naphthyl;
 X is CR₁ or N;
 Y is CR₂;
 Z is N; and
 R₁ and R₂ are each hydrogen.

[0020] Advantageously, the present invention is particularly useful for the preparation of 5,7-dihalo-6-aryl-1,2,4-triazolo[1,5-a]pyrimidines of formula I wherein

X₁ is chlorine;
 R is phenyl optionally substituted with one or more halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄-alkoxy or C₁-C₄haloalkoxy groups;
 X and Z are N; and
 Y is CH.


[0021] The process of the present invention can produce surprisingly high yields of dihydroxyazolopyrimidines and dihaloazolopyrimidines. One key factor is the temperature of the reaction between the malonic acid ester and the heterocyclamine and the use of a base. A solvent may also enhance the yield in some embodiments. Those skilled in the art will be able, without undue experimentation, to select a favorable combination of temperature and optional base and/or solvent for any particular embodiment within the scope of this invention, upon consideration of the foregoing description of the preferred embodiments and the Examples that follow.

[0022] In order to facilitate a further understanding of the invention, the following illustrative examples are presented.

Example 1

Preparation of 5,7-Dihydroxy-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine, 3-amino-1,2,4-triazole salt

[0023]



(Comparative Example)

[0024] A mixture of diethyl (2-chloro-6-fluorophenyl)-malonate (29 g, 0.1 mol), 3-amino-1,2,4-triazole (8.4 g, 0.1 mol), and the solvent mesitylene (10 mL) is heated at 160°C for 7 hours and filtered to obtain a solid. The solid is washed with diisopropyl ether and dried to give the title product as a solid (18 g, 50 % yield, mp 260-266°C).

(According to the invention)


[0025] Following essentially the same procedure, but using the appropriate solvent and/or base, the 5,7-dihydroxy-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine salts shown in Table I are obtained.

TABLE I

Solvent	Base	Temperature °C	% yield	salt
mesitylene	triethylamine	160	32	triethylamine
toluene	triethylamine	170	64	triethylamine
no added solvent	triethylamine	160	64	triethylamine
no added solvent	quinoline	180	20	quinoline

EXAMPLE 2**Preparation of 5,7-Dichloro-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine**


[0026]



[0027] A mixture of 5,7-dihydroxy-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine, 3-amino-1,2,4-triazole salt (34.8 g, 0.095 mol), and phosphorus oxychloride (100 mL) is heated in an autoclave at 140°C (2.8 bar) for 4 hours and excess phosphorus oxychloride is removed by distillation. The resultant reaction mixture is cooled to room temperature and poured into a water/dichloromethane mixture (300 mL, 1:1) while maintaining the temperature of the mixture below 30°C. The organic phase is separated, dried over anhydrous sodium sulfate, and concentrated in vacuo to obtain an oil which crystallizes overnight to give the title product as a solid (22.4 g, 74% yield, mp 118-120°C).

EXAMPLE 3**Preparation of 5,7-Dichloro-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine**


[0028]



[0029] A mixture of 3-amino-1,2,4-triazole (12.6 g, 0.15 mol), diethyl (2-chloro-6-fluorophenyl)malonate (47.6 g, 0.15 mol), and tributyl amine (27.8 g, 0.15 mol) is heated at 170°C while allowing ethanol generated during the reaction to distill off. After 2 hours, residual ethanol is removed with a slow nitrogen stream for 30 minutes. The reaction mixture is then cooled to 130°C and phosphorus oxychloride (69 g, 0.45 mol) is added dropwise over 20 minutes. The resultant clear, brown solution is refluxed for 6 hours, cooled to room temperature, and slowly added to a toluene/water (5:6) mixture (1,100 mL) with stirring. The organic phase is separated, washed sequentially with dilute hydrochloric acid and water, dried over anhydrous sodium sulfate and concentrated in vacuo to give a brown, viscous oil (44.5 g) which contains 90% of the title product (83% yield).

EXAMPLE 4**Preparation of 5,7-Dihydroxy-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine -**

[0030]



[0031] A mixture of diethyl (2-chloro-6-fluorophenyl)-malonate (7.3 g, 0.025 mol), 3-amino-1,2,4-triazole (2.1 g, 0.025 mol), mesitylene (20 mL), and pyridine (5 mL) is refluxed for 7 hours at 170°C, cooled to room temperature, and decanted to obtain a solid. A solution of the solid in water (50 mL) is acidified with concentrated hydrochloric acid (5 mL), and the resultant precipitate is collected, washed with water, and dried to give the title product as a solid (5 g, 71% yield, mp 220°C).


[0032] Following essentially the same procedure, but using the appropriate solvent and/or base, 5,7-dihydroxy-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine is obtained in the yields shown in Table II.

TABLE II

<u>Solvent</u>	<u>Base</u>	<u>Temperature °C</u>	<u>% Yield</u>
mesitylene	sodium hydroxide	170	27
mesitylene	potassium tert-butoxide	170	28
mesitylene	4-(N,N-dimethylamino)pyridine	150	61
mesitylene	quinoline	180	48
mesitylene	sodium ethoxide	170	55
SHELLSOL®	pyridine	180	38
no added solvent	pyridine	160	42
no added solvent	N,N,N',N'-tetramethylurea	170	50

COMPARATIVE EXAMPLE**Preparation of 5,7-Dihydroxy-6-(2-chloro-6-fluorophenyl)-1,2,4-triazolo[1,5-a]pyrimidine**


[0033]



[0034] Diethyl (2-chloro-6-fluorophenyl)malonate (108 g, 0.37 mol) and 3-amino-1,2,4-triazole (31.2 g, 0.37 mol) are added to a sodium ethoxide solution (previously prepared by dissolving sodium (8.5 g, 0.37 mol) in ethanol (250 mL)). The resultant reaction mixture is refluxed for 50 hours, cooled to room temperature and filtered to obtain a solid which is washed with diisopropyl ether. A solution of the washed solid in water is acidified with concentrated hydrochloric acid, and the resultant precipitate is collected, washed with water and dried to give the title product as a solid (15.7 g, 14.5% yield, mp 215°C).

Claims

1. A process for the preparation of a compound having the structural formula I



wherein

X¹ is chlorine or bromine;

R is phenyl optionally substituted with one or more halogen, nitro, cyano, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₄alkoxycarbonyl, phenyl, phenoxy or benzyloxy groups, naphthyl optionally substituted with one or more halogen, nitro, cyano, C₁-C₆alkyl,

C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy,
 C_1 - C_6 -haloalkoxy, C_1 - C_4 alkoxycarbonyl, phenyl, phenoxy or benzyloxy groups,
 C_1 - C_6 alkyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_4 alkyl,
 C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy or C_1 - C_4 haloalkoxy groups,
 C_3 - C_8 cycloalkyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 -
haloalkyl,
 C_1 - C_4 alkoxy or C_1 - C_4 haloalkoxy groups, or
 C_2 - C_6 alkenyl optionally substituted with one or more halogen, nitro, cyano, C_1 - C_4 alkyl,
 C_1 - C_4 haloalkyl,
 C_1 - C_4 alkoxy or C_1 - C_4 haloalkoxy groups;


X is CR_1 or N;

Y is CR_2 or N;

Z is CR_3 or N;

R^1 , R^2 and R^3 are each independently hydrogen or C_1 - C_6 alkyl optionally substituted with one or more halogen,
nitro, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, amino, C_1 - C_4 alkylamino
or di(C_1 - C_4 alkyl)amino groups;
which process comprises

(a) reacting (1) a malonic acid ester having the structural formula II




wherein R^8 and R^9 are each independently C_1 - C_6 alkyl, and R is as described above with
 (2) a heterocyclamine having the structural formula III



wherein X, Y and Z are as described above at a temperature of at least 100°C in the presence
 of at least one molar equivalent, relative to the malonic acid ester, of a base to form an
 intermediate salt,

(b) optionally acidifying said intermediate salt with aqueous acid to form a dihydroxyazol-
 opyrimidine having the structural formula IV



10 wherein R, X, Y and Z are as described above, and

(c) halogenating the intermediate salt or dihydroxyazolopyrimidine with at least two molar equivalents of a halogenating agent.

- 15
2. The process according to claim 1 wherein said base is selected from the group consisting of a tertiary amine, an alkali metal hydroxide, an alkaline earth metal hydroxide, an alkali metal C₁-C₆alkoxide, an alkaline earth metal C₁-C₆alkoxide, an alkali metal carbonate, and an alkaline earth metal carbonate.

20

 3. The process according to claim 2 wherein said tertiary amine is selected from the group consisting of a tri (C₂-C₆alkyl)-amine, pyridine, a substituted pyridine, quinoline, a substituted quinoline, and N,N,N',N'-tetramethylurea.

25

 4. The process according to claim 1 wherein said halogenating agent is selected from the group consisting of phosphorus oxychloride, phosphorus oxybromide, phosphorus pentachloride, phosphorus pentabromide and a suitable mixture thereof, and wherein said halogenating step is performed at a temperature of at least 100°C.

30

 5. The process according to claim 1 wherein said malonic acid ester is reacted with said heterocyclamine at a temperature of 120°C to 200°C.

35

 6. The process according to claim 1 wherein said malonic acid ester is reacted with said heterocyclamine in the presence of a solvent.

40

 7. The process according to claim 6 wherein said solvent has a boiling point of 80°C to 220°C.

45

 8. The process according to claim 6 wherein said solvent is selected from the group consisting of an aromatic hydrocarbon, a chlorinated aromatic hydrocarbon, a polynuclear aromatic hydrocarbon, an alcohol, and mixtures thereof, and the boiling point of the solvent is at least 80°C.

50

 9. The process according to claim 8 wherein said aromatic hydrocarbon is selected from the group consisting of mesitylene, toluene, a xylene, and mixtures thereof, said polynuclear aromatic hydrocarbon is selected from the group consisting of naphthalene, an alkylnaphthalene, and mixtures thereof, and said alcohol is butanol.

55

 10. The process according to claim 1 wherein said aqueous acid is an aqueous mineral acid selected from the group consisting of hydrochloric acid, hydrobromic acid, and sulfuric acid.
 11. The process according to claim 1 wherein said halogenation is conducted at a pressure greater than one atmosphere.
 12. The process according to claim 1 wherein

X¹ is chlorine;

R is phenyl optionally substituted with one or more halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, phenyl, phenoxy or benzyloxy groups, or naphthyl;

X is CR₁ or N;

Y is CR₂;

Z is N; and

5 R¹ and R² are each hydrogen.

13. A process for the preparation of a compound of formula IV according to claim 1, which process comprises reaction steps (a) and (b) according to claim 1.

10 14. The process according to claim 13 wherein in formula IV

R is phenyl optionally substituted with one or more halogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, phenyl, phenoxy or benzyloxy groups, or naphthyl;

15 X is CR₁ or N;

Y is CR₂;

Z is N; and


20 R¹ and R² are each hydrogen.

Patentansprüche

25

1. Verfahren zur Herstellung einer Verbindung mit der Strukturformel I

30



(I)

35

in der

40 X¹ Chlor oder Brom bedeutet,

R Phenyl, das gegebenenfalls durch eine oder mehrere Halogen-, Nitro-, Cyan-, C₁-C₆-Alkyl-, C₁-C₆-Halogenalkyl-, C₁-C₆-Alkoxy-, C₁-C₆-Halogenalkoxy-, C₁-C₄-Alkoxy-carbonyl-, Phenyl-, Phenoxy- oder Benzyloxygruppen substituiert ist, Naphthyl, das gegebenenfalls durch eine oder mehrere Halogen-, Nitro-, Cyan-, C₁-C₆-Alkyl-, C₁-C₆-Halogenalkyl-, C₁-C₆-Alkoxy-, C₁-C₆-Halogenalkoxy-, C₁-C₄-Alkoxy-carbonyl-, Phenyl-, Phenoxy- oder Benzyloxygruppen substituiert ist, C₁-C₆-Alkyl, das gegebenenfalls durch eine oder mehrere Halogen-, Nitro-, Cyan-, C₁-C₄-Alkyl-, C₁-C₄-Halogenalkyl-, C₁-C₄-Alkoxy- oder C₁-C₄-Halogenalkoxygruppen substituiert ist, C₃-C₈-Cycloalkyl, das gegebenenfalls durch eine oder mehrere Halogen-, Nitro-, Cyan-, C₁-C₄-Alkyl-, C₁-C₄-Halogenalkyl-, C₁-C₄-Alkoxy- oder C₁-C₄-Halogenalkoxygruppen substituiert ist, oder C₂-C₆-Alkenyl, das gegebenenfalls durch eine oder mehrere Halogen-, Nitro-, Cyan-, C₁-C₄-Alkyl-, C₁-C₄-Halogenalkyl-, C₁-C₄-Alkoxy- oder C₁-C₄-Halogenalkoxygruppen substituiert ist, bedeutet,

55 X CR₁ oder N bedeutet,


Y CR₂ oder N bedeutet,

Z CR₃ oder N bedeutet,

R¹, R² und R³ jeweils unabhängig Wasserstoff oder C₁-C₆-Alkyl, das gegebenenfalls durch eine oder mehrere Halogen-, Nitro-, Cyan-, C₁-C₄-Alkyl-, C₁-C₄-Halogenalkyl-, C₁-C₄-Alkoxy-, C₁-C₄-Halogenalkoxy-, Amino-, C₁-C₄-Alkylamino- oder Di(C₁-C₄-alkyl)aminogruppen substituiert ist, bedeuten,

dadurch gekennzeichnet, daß man

(a) (1) einen Malonsäureester mit der Strukturformel II




in der R⁸ und R⁹ jeweils unabhängig C₁-C₆-Alkyl bedeuten und R die oben beschriebene Bedeutung aufweist, mit (2) einem Heterocyclamin mit der Strukturformel III



in der X, Y und Z die oben beschriebene Bedeutung aufweisen, bei einer Temperatur von mindestens 100°C in Gegenwart mindestens eines Moläquivalents (in bezug auf den Malonsäureester) einer Base umsetzt, wodurch man ein Salz als Zwischenprodukt erhält,

(b) gegebenenfalls dieses als Zwischenprodukt erhaltene Salz mit wäßriger Säure ansäuert, wodurch man ein Dihydroxyazolopyrimidin mit der Strukturformel IV



in der R, X, Y und Z die oben beschriebene Bedeutung aufweisen, erhält, sowie

(c) das als Zwischenprodukt erhaltene Salz bzw. Dihydroxyazolopyrimidin mit mindestens zwei Moläquivalenten eines Halogenierungsmittels halogeniert.


2. Verfahren nach Anspruch 1, wobei die Base aus der Gruppe tertiäres Amin, Alkalimetallhydroxid, Erdalkalimetallhydroxid, Alkalimetall-C₁-C₆-alkoholat, Erdalkalimetall-C₁-C₆-alkoholat, Alkalimetallcarbonat und Erdalkalimetallcarbonat stammt.
3. Verfahren nach Anspruch 2, wobei das tertiäre Amin aus der Gruppe Tri(C₂-C₆-alkyl)amin, Pyridin, substituiertes

Pyridin, Chinolin, substituiertes Chinolin und N,N,N',N'-Tetramethylharnstoff stammt.

- 5 4. Verfahren nach Anspruch 1, wobei das Halogenierungsmittel aus der Gruppe Phosphoroxychlorid, Phosphorox-
bromid, Phosphorpentachlorid, Phosphorpentabromid sowie eine geeignete Mischung aus diesen Substanzen
stammt und wobei der Halogenierungsschritt bei einer Temperatur von mindestens 100°C durchgeführt wird.
- 5 5. Verfahren nach Anspruch 1, wobei der Malonsäureester mit dem Heterocyclylamin bei einer Temperatur von 120°C
bis 200°C umgesetzt wird.
- 10 6. Verfahren nach Anspruch 1, wobei der Malonsäureester mit dem Heterocyclylamin in Gegenwart eines Lösungs-
mittels umgesetzt wird.
7. Verfahren nach Anspruch 6, wobei das Lösungsmittel einen Siedepunkt von 80°C bis 220°C aufweist.
- 15 8. Verfahren nach Anspruch 6, wobei das Lösungsmittel aus der Gruppe aromatischer Kohlenwasserstoff, chlorierter
aromatischer Kohlenwasserstoff, mehrkerniger aromatischer Kohlenwasserstoff, Alkohol und deren Mischungen
stammt und der Siedepunkt des Lösungsmittels mindestens 80°C beträgt.
- 20 9. Verfahren nach Anspruch 8, wobei der aromatische Kohlenwasserstoff aus der Gruppe Mesitylen, Toluol, Xylol
und deren Mischungen stammt, der mehrkernige aromatische Kohlenwasserstoff aus der Gruppe Naphthalin, Al-
kyl-naphthalin und deren Mischungen stammt und es sich bei dem Alkohol um Butanol handelt.
- 25 10. Verfahren nach Anspruch 1, wobei es sich bei der wäßrigen Säure um eine wäßrige Mineralsäure aus der Gruppe
Chlorwasserstoffsäure, Bromwasserstoffsäure und Schwefelsäure handelt.
11. Verfahren nach Anspruch 1, wobei die Halogenierung bei einem Druck von mehr als einer Atmosphäre durchge-
führt wird.
- 30 12. Verfahren nach Anspruch 1, wobei
 - X¹ Chlor bedeutet,
 - R Phenyl, das gegebenenfalls durch eine oder mehrere Halogen-, C₁-C₄-Alkyl-, C₁-C₄-Halogenalkyl-,
C₁-C₄-Alkoxy-, C₁-C₄-Halogenalkoxy-, Phenyl-, Phenoxy- oder Benzyloxygruppen substituiert ist,
oder Naphthyl bedeutet,
 - 35 X CR₁ oder N bedeutet,
 - Y CR₂ bedeutet,
 - 40 Z N bedeutet und
 - R¹ und R² jeweils Wasserstoff bedeuten.
- 45 13. Verfahren zur Herstellung einer Verbindung der Formel IV nach Anspruch 1, das Reaktionsschritte (a) und (b)
nach Anspruch 1 umfaßt.
- 50 14. Verfahren nach Anspruch 13, wobei, in Formel IV,
R Phenyl, das gegebenenfalls durch eine oder mehrere Halogen-, C₁-C₄-Alkyl-, C₁-C₄-Halogenalkyl-, C₁-C₄-Alk-
oxy-, C₁-C₄-Halogenalkoxy-, Phenyl-, Phenoxy- oder Benzyloxygruppen substituiert ist, oder Naphthyl bedeutet,
 - X CR₁ oder N bedeutet,
 - Y CR₂ bedeutet,
 - 55 Z N bedeutet und
 - R¹ und R² jeweils Wasserstoff bedeuten.

Revendications

1. Procédé pour la préparation d'un composé répondant à la formule développée I



dans laquelle

X¹ représente un atome de chlore ou un atome de brome ;

R représente un groupe phényle, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, nitro, cyano, alkyle en C₁ - C₆, halogénaalkyle en C₁ - C₆, alcoxy en C₁ - C₆, halogénalcoxy en C₁ - C₆, alcoxy(en C₁ - C₄)carbonyle, phényle, phénoxy ou benzyloxy, un groupe naphthyle, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, nitro, cyano, alkyle en C₁ - C₆, halogénaalkyle en C₁ - C₆, alcoxy en C₁ - C₆, halogénalcoxy en C₁ - C₆, alcoxy(en C₁ - C₄)carbonyle, phényle, phénoxy ou benzyloxy, un groupe alkyle en C₁ - C₆, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, nitro, cyano, alkyle en C₁ - C₄, halogénaalkyle en C₁ - C₄, alcoxy en C₁ - C₄ ou halogénalcoxy en C₁ - C₄, un groupe cycloalkyle en C₃ - C₈, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, nitro, cyano, alkyle en C₁ - C₄, halogénaalkyle en C₁ - C₄, alcoxy en C₁ - C₄ ou halogénalcoxy en C₁ - C₄, ou un groupe alcényle en C₂ - C₆, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, nitro, cyano, alkyle en C₁ - C₄, halogénaalkyle en C₁ - C₄, alcoxy en C₁ - C₄ ou halogénalcoxy en C₁ - C₄ ;

X représente un groupe CR¹ ou un atome d'azote ;


Y représente un groupe CR² ou un atome d'azote ;

Z représente un groupe CR³ ou un atome d'azote ;

R¹, R² et R³ représentent, chacun indépendamment l'un de l'autre, un atome d'hydrogène ou un groupe alkyle en C₁ - C₆, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, nitro, cyano, alkyle en C₁ - C₄, halogénaalkyle en C₁ - C₄, alcoxy en C₁ - C₄, halogénalcoxy en C₁ - C₄, amino, alkyl(en C₁ - C₄)amino ou dialkyl(en C₁ - C₄)amino ;

ledit procédé comprenant le fait de

(a) faire réagir (1) un ester d'acide malonique répondant à la formule développée II




dans laquelle R⁸ et R⁹ représentent, chacun indépendamment l'un de l'autre, un groupe alkyle en C₁ - C₆, et R est tel que décrit ci-dessus avec (2) une hétérocyclamine répondant à la formule développée III



dans laquelle X, Y et Z sont tels que décrits ci-dessus, à une température d'au moins 100 °C en présence d'au moins un équivalent molaire, par rapport à l'ester de l'acide malonique, d'une base pour former un sel intermédiaire,

(b) le cas échéant, acidifier ledit sel intermédiaire avec un acide aqueux pour former une dihydroxyazolopyrimidine répondant à la formule développée IV



dans laquelle R, X, Y et Z sont tels que décrits ci-dessus, et

(c) soumettre à une halogénéation le sel intermédiaire ou la dihydroxyazolopyrimidine avec au moins deux équivalents molaires d'un agent d'halogénéation.

2. Procédé selon la revendication 1, dans lequel ladite base est choisie parmi le groupe constitué par une amine tertiaire, un hydroxyde de métal alcalin, un hydroxyde de métal alcalino-terreux, un alcoxyde en C₁ - C₆ de métal alcalin, un alcoxyde en C₁ - C₆ de métal alcalino-terreux, un carbonate de métal alcalin et un carbonate de métal alcalino-terreux.
3. Procédé selon la revendication 2, dans lequel ladite amine tertiaire est choisie parmi le groupe constitué par une tri(alkyl(en C₂ - C₆))amine, de la pyridine, une pyridine substituée, de la quinoléine, une quinoléine substituée et de la N,N,N',N'-tétraméthylurée.
4. Procédé selon la revendication 1, dans lequel ledit agent d'halogénéation est choisi parmi le groupe constitué par l'oxychlorure de phosphore, l'oxybromure de phosphore, le pentachlorure de phosphore, le pentabromure de phosphore et un de leurs mélanges appropriés, et dans lequel ladite étape d'halogénéation est effectuée à une température d'au moins 100 °C.
5. Procédé selon la revendication 1, dans lequel on fait réagir ledit ester de l'acide malonique avec ladite hétérocyclamine à une température de 120 °C à 200 °C.
6. Procédé selon la revendication 1, dans lequel on fait réagir ledit ester de l'acide malonique avec ladite hétérocyclamine en présence d'un solvant.
7. Procédé selon la revendication 6, dans lequel ledit solvant possède un point d'ébullition de 80 °C à 220 °C.
8. Procédé selon la revendication 6, dans lequel ledit solvant est choisi parmi le groupe constitué par un hydrocarbure aromatique, un hydrocarbure aromatique chloré, un hydrocarbure aromatique polynucléaire, un alcool et leurs mélanges, et le point d'ébullition du solvant s'élève à au moins 80 °C.
9. Procédé selon la revendication 8, dans lequel ledit hydrocarbure aromatique est choisi parmi le groupe constitué

par le mésitylène, le toluène, un xylène et leurs mélanges, ledit hydrocarbure aromatique polynucléaire est choisi parmi le groupe constitué par le naphthalène, un alkylnaphtalène et leurs mélanges et ledit alcool est le butanol.

10. Procédé selon la revendication 1, dans lequel ledit acide aqueux est un acide minéral aqueux choisi parmi le groupe constitué par l'acide chlorhydrique, l'acide bromhydrique et l'acide sulfurique.

11. Procédé selon la revendication 1, dans lequel ladite halogénéation est effectuée à une pression supérieure à une atmosphère.

12. Procédé selon la revendication 1, dans lequel

X¹ représente un atome de chlore ;

R représente un groupe phényle, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, alkyle en C₁ - C₄, halogénalkyle en C₁ - C₄, alcoxy en C₁ - C₄, halogénalcoxy en C₁ - C₄, phényle, phénoxy ou benzyloxy, ou représente un groupe naphthyle ;

X représente un groupe CR¹ ou un atome d'azote ;

Y représente un groupe CR² ;

Z représente un atome d'azote ; et

R¹ et R² représentent chacun un atome d'hydrogène.

13. Procédé pour la préparation d'un composé répondant à la formule IV selon la revendication 1, ledit procédé comprenant les étapes réactionnelles (a) et (b) selon la revendication 1.

14. Procédé selon la revendication 13, dans lequel, dans la formule IV

R représente un groupe phényle, portant le cas échéant à titre de substituants un ou plusieurs groupes halogéno, alkyle en C₁ - C₄, halogénalkyle en C₁ - C₄, alcoxy en C₁ - C₄, halogénalcoxy en C₁ - C₄, phényle, phénoxy ou benzyloxy ; ou représente un groupe naphthyle ;

X représente un groupe CR¹ ou un atome d'azote ;

Y représente un groupe CR² ;

Z représente un atome d'azote ; et

R¹ et R² représentent chacun un atome d'hydrogène.